

Potato Late Blight Fungicides Registered for WI, 2024.

In-furrow and seed treatment registrations are omitted. Not a comprehensive list. Most fungicides listed are for use in conventional systems. Where generic fungicide trade names are listed, there may be numerous. Compiled 19 May 2024.

Amanda J. Gevens, Extension Plant Pathologist, UW-Madison

Cell Phone: 608-575-3029; E-mail: gevens@wisc.edu

Trade Name (rate/A)	Active Ingredient(s)	PHI	REI	FRAC #	Activity of Fungicide	Comments
<u>Agri Tin, Super Tin 4L, Super Tin 80WP</u> (4-6 fl oz)	triphenyltin hydroxide	7 days	48 hours	30	Protectant; kills spores on contact	Restricted use pesticide. 3 fl oz rate can be used if material is tank-mixed with another fungicide.
<u>Alude</u> (1.25 qt in 90 gal water) <u>Confine Extra</u> (3-5 qt in 20 gal water/acre) <u>K-Phite 7LP</u> (1-3 qt in 10 gal water/acre)	mono and dipotassium salts of phosphorous acid	0 days	4 hours	33	Upregulates resistance/disease protection in plant	Foliar application. Use higher rate when disease pressure is severe. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or that are heat or moisture stressed.
<u>Ariston</u> (2 pt)	chlorothalonil + cymoxanil	14 day	12 hours	M5+2 7	Protectant and locally systemic	Additional chlorothalonil may be tank-mixed with this formulation to enhance % active ingredient applied (be sure to include the Ariston component in overall season total). Cymoxanil is Curzate.
<u>Elixir</u> (1.2-2.0 lb)	mancozeb+ chlorothalonil	7 days	24 hours	M3+ M5	Protectant	Use higher rate as vines increase in size.
<u>Fosphite, Rampart</u> (1-4 qt)	potassium phosphite	0 days	4 hours	33	Upregulates resistance/disease protection in plant	Foliar post-emergence spray and post harvest spray for control in storage. Apply in at least 20 gal water/acre.
<u>Fungi-Phite</u> (Foliar: 2 qt/A Seed trt: 15% vol to vol-2 ton in 1 gal solution) <u>Helena Prophyt</u> (4 pt/acre) <u>Badge SC</u> (1-3 pt)	potassium phosphite	0 days	4 hours	33	Upregulates resistance/disease protection in plant	Seed piece spray and foliar post-emergence spray. Tank-mix with another effective fungicide is recommended and use high label rate for late blight control.
<u>Bravo Ultrex</u> (.7 then .9 to 1.36 lb) <u>Bravo WeatherStik, Echo 720, Equus 720 SST, Initiate 720, Chlorothalonil 720 SC, Chloronil 720, Praiz</u> (.75 then 1-1.5 pt) <u>Bravo Zn, Equus 500 Zn, Initiate Zn</u> (1 1/8 then 1.5 to 2.25 pt)	copper hydroxide, copper oxychloride	0 days	24 hours	M1	Protectant	Protectant activity only. Apply at 7 to 10 day interval.
<u>Bravo Ultrex</u> (.7 then .9 to 1.36 lb) <u>Bravo WeatherStik, Echo 720, Equus 720 SST, Initiate 720, Chlorothalonil 720 SC, Chloronil 720, Praiz</u> (.75 then 1-1.5 pt) <u>Bravo Zn, Equus 500 Zn, Initiate Zn</u> (1 1/8 then 1.5 to 2.25 pt)	chlorothalonil	7 days	12 hours	M5	Protectant	11.25 lb a.i./acre maximum on standard label. However, WI has a special 24(c) registration for long season potatoes extending the max a.i. from 11.25 to 16 lb a.i./acre with Bravo (Syngenta) and Echo (Sipcam Advan) formulations.

Potato Late Blight Fungicides Registered for WI, 2024.

Trade Name (rate/A)	Active Ingredient(s)	PHI	REI	FRAC #	Activity of Fungicide	Comments
<p><u>Echo Zn</u> (1 to 2.125 pt)</p> <p><u>Equus DF</u> (.7 then .9 to 1.36 lb)</p> <p>Echo 90DF (5/8 then 7/8 to 1.25 lb)</p>	chlorothalonil	7 days	12 hours	M5	Protectant	11.25 lb a.i./acre maximum on standard label. However, WI has a special 24(c) registration for long season potatoes extending the max a.i. from 11.25 to 16 lb a.i./acre with Bravo (Syngenta) and Echo (Sipcam Advan) formulations.
<u>Cabrio Plus</u> (2.9 lb)	pyraclostrobin + metiram	3 days	24 hours	11+M3	Locally systemic and protectant	17.4 lb/acre maximum per season. Do not apply more than 2 sequential applications.
<p><u>Champ WG</u> (1 to 1.5 lb 3 to 4 lb in severe areas)</p> <p><u>Champ Formula 2 Flowable</u> (2/3 to 2 2/3 pt)</p> <p><u>Champ DP Dry Prill</u> (2/3 to 1 lb 2 to 2 2/3 lb when disease is severe)</p> <p><u>Kentan DF</u> (1-2.5 lb 4 lb when severe)</p> <p><u>Kocide 2000, Kocide 3000</u> (.73- 3 lb .5-1.75 lb)</p> <p><u>Nu-Cop 3L</u> (2/3 to 2 pt 2 to 4 pt if severe)</p> <p><u>Nu-Cop 50DF</u> (1-1.5 lb 3-4 lb if severe)</p> <p><u>Previsto</u> (1-3 qt)</p>	copper hydroxide	0 days	24 hours	M1	Protectant	Use high label rates for foliar late blight protection.
<p><u>C-O-C-S WDG</u> (1.5- 4 lb)</p> <p><u>Cuprofix-Ultra 40 Dispers</u> (0.75-3.0 lb)</p>	copper oxychloride, basic copper sulfate	0 days	24 hours	M1	Protectant	Use high label rates for foliar late blight protection.
<u>Mastercop</u> (0.5-1.5 pt)	copper sulfate pentahydrate	0 days	24 hours	M1	Protectant	Use high label rates for foliar late blight protection.

Potato Late Blight Fungicides Registered for WI, 2024.

Trade Name (rate/A)	Active Ingredient(s)	PHI	REI	FRAC #	Activity of Fungicide	Comments
<u>Cueva</u> (2 gal in 50-100 gal water/acre)	copper octanoate	0 days	24 hours	M1	Protectant	Use high label rates for foliar late blight protection.
<u>Curzate 60DF</u> (3.2 oz foliar)	cymoxanil	14 days	12 hours	27	Locally systemic	Locally-systemic fungicide. Must be tank-mixed with a protectant fungicide. Rainfast within 2 hours. Cymbol.
<u>Dithane F45 Rainshield</u> (.4 to 1.6 qt) <u>Dithane M45</u> (.5 to 2 lb) <u>Koverall,</u> <u>Roper DF</u> <u>Rainshield,</u> <u>Fortuna</u> <u>75WDG</u> (1-2.0 lb)	mancozeb	24 hours	3 days	M3	Protectant	Max rate per acre/season is 11.2 lb a.i. Begin use at lower rate and increase as vines increase in size.
<u>Evito 480SC,</u> <u>Aftershock</u> (3.8 fl oz)	fluoxastrobin	7 days	12 hours	11	Locally systemic	Follow label for resistance management.
<u>Forum</u> (Foliar and tuber control: 6 oz)	dimethomorph	4 days	12 hours	40	Systemic	May be tank-mixed with another effective fungicide for enhanced management – but not required by label. Addition of an adjuvant may enhance management. Can be applied after vine kill.
<u>Gavel 75DF</u> (1.5 to 2 lb)	zoxamide+ mancozeb	3 days	48 hours	22+M3	Protectant	Do not make >6 applications/crop. Contact fungicide.
<u>Gem 500SC</u> (3.8 fl oz)	trifloxystrobin	7 days	12 hours	11	Locally systemic	Follow label for resistance management.
<u>Headline</u> (6 to 12 fl oz)	pyraclostrobin	3 days	12 hours	11	Locally systemic	Follow label for resistance management.
<u>ManKocide</u> (1.5 to 2 then 4-5 lb)	mancozeb+ copper hydroxide	3 days	24 hours	M3+ M1	Protectant	Not labeled as a seed trt for potatoes.
<u>Omega 500F</u> (5.5 fl oz)	fluazinam	14 days	48 hours	29	Protectant	REI is 4 days for high exposure activities. New special local need label 24c in April 2011.
<u>Omega Top MP</u> (5.5 fl oz) – individual label for Omega sold in co-pack with Top MP (difenoconazole)	fluazinam	14 days	48 hours	29	Protectant and locally systemic	Can be applied aerially. REI is 4 days for high exposure activities.

Potato Late Blight Fungicides Registered for WI, 2024.

Trade Name (rate/A)	Active Ingredient(s)	PHI	REI	FRAC #	Activity of Fungicide	Comments
<u>Orondis Ultra</u> (5.5-8.0 fl oz)	oxathiapiprolin + mandipropamid	14 days	4 hours	U15 + 40	Translaminar or systemic within leaf and xylem mobile; rainfast within 30 minutes	Max single application rate is 8.0 fl oz/A. Max annual rate is 32.0 fl oz/A/year. Do not apply >0.125 lb ai/year of oxathiapiprolin. Do not apply >0.522 lb ai/year of mandipropamid. Make no more than 2 sequential applications before rotating to a different mode of action. Do not follow soil applications of oxathiapiprolin with foliar applications of Orondis Ultra.
<u>Orondis Opti</u> (1.75-2.50 pt)	oxathiapiprolin + chlorothalonil	7 days	12 hours	U15 + M5	Translaminar or systemic within leaf and xylem mobile; rainfast within 30 minutes	Max single application rate is 2.5 pt/A. Max application annual rate is 10 pt/A/year. Do not apply >0.125 lb ai/A/year of oxathiapiprolin. Do not apply >11.25 lb ai/A/year of chlorothalonil.
<u>Oxidate</u> (40 to 120 fl oz to 100 gal water, 30-100 gal/acre)	hydrogen dioxide	0 days	1 hour	NC	Kills spores on contact and no residual protectant activity after treatment	Foliar spray for late blight. Frequent applications (5-day intervals) can limit sporulation.
<u>Penncozeb 80WP</u> , <u>Penncozeb 75DF</u> (.5 to 2 lb) <u>Penncozeb 4FL</u> , <u>Manzate flowable</u> (.4 to 1.6 qt) <u>Manzate Pro-Stick</u> (1 to 2 lb, seed trt: 1.25 lb/50 gal water)	mancozeb	3 days	24 hours	M3	Protectant	Do not exceed 11.2 lb a.i./acre/year.
<u>Phostrol</u> (2.5 to 10 pt) (Post harvest trt: 1 gal/ton in .5 gal water)	mono- and di-basic sodium, potassium, and ammonium phosphites	0 days	4 hours	33	Upregulates resistance or plant defense	Can be applied as a foliar for late blight, pink rot, and Pythium leak. Can be applied post-harvest for storage disease control.
<u>Polyram 80DF</u> (1.5 to 2 lb in 15 gal water/acre minimum)	metiram	3 days	24 hours	M3	Protectant	Metiram is an EBDC, like mancozeb (M3). Total amount of a.i. per year/acre must include all EBDCs.
<u>Previcur Flex</u> (.7 to 1.2 pt)	propamocarb hydrochloride	14 days	12 hours	F	Systemic antisporeulant	Apply in a tank-mix with effective protectant. Can be applied as a broadcast or banded application over the row, post-emergence.
<u>Priaxor</u> (4-8 fl oz)	fluxapyroxad+ pyraclostrobin	7 days	12 hours	7+11	Protectant and locally systemic	Cannot apply more than 3 applications/season. Follow label for resistance management. Xemium and Headline pre-mix.

Potato Late Blight Fungicides Registered for WI, 2024

Trade Name (rate/A)	Active Ingredient(s)	PHI	REI	FRA C #	Activity of Fungicide	Comments
<u>Quadris</u> , <u>Satori</u> , <u>Willowood</u> <u>Azoxy 2SC</u> , <u>Aframe</u> , <u>Equation SC</u> (6 to 15.5 fl oz) <u>Trevo</u> , <u>Azoxystrar</u> , <u>Azoxyzone</u> (6-20 fl oz)	azoxystrobin	14 days	4 hours	11	Locally systemic	Alternate away from Group 11 fungicides to manage resistance.
<u>Quadris Opti</u> (1.6 pt)	azoxystrobin+chlorothalonil	14 days	12 hours	11+M5	Locally systemic and protectant	Alternate away from Group 11 fungicides to manage resistance.
<u>Ranman</u> (1.4 to 2.75 fl oz)	cyazofamid	7 days	12 hours	21	Protectant	Follow label for resistance management.
<u>Reason</u> (5.5 to 8.2 fl oz)	fenamidone	14 days	12 hours	11	Locally systemic	Follow label for resistance management.
<u>Revus Top</u> (5.5 to 7 fl oz)	mandipropamid+difenoconazole	14 days	12 hours	40+3	Locally systemic and contact	Addition of an adjuvant is recommended.
<u>Tanos</u> (8 to 10 oz)	cymoxanil + famoxadone	14 days	12 hours	27+11	Locally systemic and contact	Must be tank-mixed with an effective protectant fungicide. Good protectant for limiting leaf blight. Excellent curative.
<u>Ridomil Gold SL</u> (1 to 2 pt)	mefenoxam	14 days	48 hours	4	Systemic	Do not apply beyond the at-planting stage.
<u>Ridomil Gold Bravo SC</u> (2.5 pt)	mefenoxam+chlorothalonil	14 days	48 hours	4+M5	Systemic and protectant	Follow label for resistance management.
<u>Ridomil Gold Copper</u> (2 lb)	mefenoxam+copper hydroxide	14 days	48 hours	4+M1	Systemic and protectant	Tank-mix with an effective protectant.
<u>Ridomil Gold MZ WG</u> (2.5 lb)	mefenoxam+mancozeb	3 days	48 hours	4+M3	Systemic and protectant	Follow label for resistance management.
<u>Zampro</u> (11-14 fl oz)	ametoctradin+dimethomorph	4 days	12 hours	45+40	Systemic and protectant	Do not make more than 2 sequential applications. Follow label for resistance management. Ametoctradin is new a.i.; dimethomorph is Forum (formerly Acrobat).
<u>Zing!</u> (32-34 fl oz)	zoxamide+chlorothalonil	7 days	12 hours	22+M5	Protectant	Do not make more than 2 sequential applications before alternating with another fungicide of a different mode of action. Do not make >8 applications or apply >1.52 lb of zoxamide and 8.88 lb of chlorothalonil per season per acre.