



Vegetable Crop Updates

*A newsletter for commercial potato and vegetable growers prepared by
University of Wisconsin-Madison vegetable research and extension specialists*

July 12, 2026

In This Issue:

- Weather Impacts on Potato & Vegetable Crops
- Potato Early and Late Blight Disease Risk Model Updates
- Cucurbit Downy Mildew Updates and Management Strategies
- Potato Virus Y Management
- Corn Earworm Management
- Langlade Co. Airport Research Station Field Day Flier

Calendar of Events:

July 16, 2026 – UW Langlade County Airport Research Station Field Day, Antigo, WI

July 23, 2026 – UW Rhinelander Agricultural Research Station Field Day, Rhinelander, WI

December 1-3, 2026 – Midwest Food Products Association Annual Convention & Expo, Processing Crops Conference, Wisconsin Dells, WI

February 9-11, 2027 – WPVGA/UWEX Grower Education Conference, Stevens Point, WI

Yi Wang, Associate Professor & Extension Potato and Vegetable Production Specialist, UW-Madison, Dept. of Plant and Agroecosystem Sciences, 608-265-4781, Email: wang52@wisc.edu.

The Wisconsin Agriculture Weather Outlook indicates that average temperatures were 4-8°F above normal across most of the state, with the eastern part of the state the most above normal. Some days had heat indices exceeding 100°F. The driest regions were the north-central and southeast counties, while some southwestern counties had rainfall totals between 3" and 6" last week.

Satellite-based soil moisture estimates indicate that drier-than-normal conditions persist across the northwest, north-central, and east-central counties. The outlook for mid-to-late July indicates statewide above-normal temperatures and below-normal precipitation probabilities. There will be minimal chances for rain over the next seven days. Late summer through early fall (July – September) might be cooler-than-normal in southern and central parts of the state, with rainfall chances leaning towards below normal for most parts of Wisconsin.



Some of the folks had questions about managing hail damage to their vegetable crops. Hail is one of the more unpredictable risks for home gardeners or commercial producers (clockwise from top left: hail damage to potatoes, tomatoes, cucumbers, and chards). There are no preventive options once a storm develops, and the damage can range from minor leaf injury to severe canopy loss and yield reduction. There are 4 factors that can determine the outcome of hail:

1. Amount of defoliation: less than 30% of defoliation often leads to minimal yield loss because there remains enough leaf area to keep photosynthesis going on. Between 30 and 70% defoliation could be associated with more yield reduction due to limited carbohydrate synthesis. Severe defoliation exceeding 70% can significantly reduce yield, especially if stems are also damaged.
2. When the hail occurs: early-season injury during crops' vegetative growth is often the least damaging because plants still have time to regrow. Tuber initiation and early tuber bulking are the most sensitive periods when plants are highly vulnerable. Defoliation during this period usually reduces tuber set and tuber size in potatoes, resulting in fewer, smaller tubers. Later in the season, during tuber maturation, hail will have less impact on yield but can influence tuber quality and increase disease risks.
3. Cultivar selection: late-maturing cultivars are better able to recover from hail damage, as they have more time in the growing season to outgrow it, particularly if favorable conditions persist. Early-maturing cultivars have a shorter recovery window, and thus similar damage at the same growth stage results in a greater proportional yield loss.
4. Weather conditions after the hail event: these can affect how well the crops recover. Warm but not high temperatures, sufficient soil moisture, and minimal biotic stress favor strong regrowth. Under these conditions, plants produce new leaves to rebuild the canopy. In comparison, hot and dry spells can compound stress and slow down recovery; in addition, prolonged wet days following hail damage can increase the risk of disease development in damaged tissues.

Foliar nutrient applications are sometimes applied to stimulate regrowth. However, there is little evidence that these nutrients translate into higher yields at harvest. For potatoes, heavy nitrogen use after hail can shift growth towards vines and away from tuber setting and bulking, leading to a higher incidence of misshapen potatoes and those with hollow heart.

In summary, the crops' natural ability to recover, in combination with favorable weather conditions, has a greater impact on yield than any input applied after the hailstorm. The best practice is to optimize recovery conditions by maintaining proper soil moisture, providing adequate but not excessive fertility, minimizing additional environmental stresses, and allowing time for regrowth. If possible, slightly extending the growing season may also help offset canopy loss by giving the plants more time to grow.

Amanda Gevens, Professor & Extension Vegetable Pathologist, UW-Madison, Dept. of Plant Pathology, 608-575-3029, gevens@wisc.edu, Lab Website: <https://vegpath.plantpath.wisc.edu/>.

Current P-Day (Early Blight) and Disease Severity Value (Late Blight) Accumulations will be posted at our website and available in the weekly newsletters. Thanks to Ben Bradford, UW- Madison Entomology for supporting this effort and providing a summary reference table: <https://agweather.cals.wisc.edu/thermal-models/potato>. A Potato Physiological Day or P-Day value of ≥ 300 indicates the threshold for early blight risk in potato and triggers preventative fungicide application. A Disease Severity Value or DSV of ≥ 18 indicates the threshold for late blight risk and triggers preventative fungicide application in potato. Data from the modeling source: <https://agweather.cals.wisc.edu/vdifn> are used to generate these risk values in the table below. I've estimated early, mid-, and late planting dates by region based on communications with stakeholders. These are intended to help in determining optimum times for preventative fungicide applications to limit early and late blight in Wisconsin.

Location	Planting Date		50% Emergence Date	Disease Severity Values (DSVs) through 7/11/26	Potato Physiological Days (P-Days) through 7/11/26
Spring Green	Early	Apr 10	May 7	25	499
	Mid	May 5	May 20	23	428
	Late	May 16	June 2	23	332
Arlington	Early	Apr 12	May 8	12	493
	Mid	May 6	May 23	12	418
	Late	May 20	June 4	12	321
Grand Marsh	Early	Apr 13	May 9	26	470
	Mid	May 6	May 22	26	411

	Late	May 21	June 5	26	306
Hancock	Early	Apr 14	May 11	21	466
	Mid	May 10	May 30	20	351
	Late	May 23	June 6	19	298
Plover	Early	Apr 15	May 12	21	465
	Mid	May 10	May 30	19	353
	Late	May 25	June 6	18	300
Antigo	Early	May 12	May 29	20	340
	Mid	May 25	June 5	21	294
	Late	June 8	June 20	12	181
Rhineland	Early	May 15	June 6	18	275
	Mid	May 28	June 10	16	238
	Late	June 9	June 25	10	143

Late blight of potato/tomato. With the exception of the Arlington area, most potato production areas have accumulated Disease Severity Values or DSVs at or surpassing the action threshold of 18 as of 7/11 – this is 0-7 more DSVs since last week. This indicates that late blight favorable weather has occurred, since ~50% crop emergence and we have accumulated enough late blight-favorable weather to warrant a preventative fungicide application to target this disease. Fields with ≥ 18 DSVs should receive preventative fungicide applications to manage for late blight.

<https://vegpath.plantpath.wisc.edu/diseases/potato-late-blight/> An updated list of fungicides for management potato late blight in Wisconsin 2026 is provided here: <https://vegpath.plantpath.wisc.edu/resources/late-blight-fungicides/>

Early blight of potato. Most potato fields south of Antigo have surpassed the treatment threshold of 300 P-Days. The earliest plantings in Antigo (emergence late May/early June) are now at or above threshold. Preventative fungicide applications are warranted for these fields. In the past week regions of the state accumulated between 57 and 64 P-Days. For support in selecting fungicides, please consult the A3422 Commercial Vegetable Production in Wisconsin: <https://vegpath.plantpath.wisc.edu/resources/a3422/>

Cucurbit Downy mildew update. No confirmations of downy mildew in cucurbits in the upper midwestern US at this time. While Dr. Mary Hausbeck's Michigan State University downy mildew spore trapping network is reporting the presence of pathogen spores in several counties of Michigan, there have been no field disease reports. Downy mildew on cucumbers and muskmelon/cantaloupe (likely Clade 2 strain type) in Georgia and Florida were reported in late June 2026. <https://cdm.ipmPIPE.org/> Based on this information, there is some risk for downy mildew on cucumbers in Wisconsin with relatively lower risk on muskmelon/cantaloupe. Even lower risk on pumpkin and squash at this time. Hotter and drier weather will reduce risk of infection and development of disease, whereas cooler and wetter weather will promote disease. For more information: <https://vegpath.plantpath.wisc.edu/diseases/cucurbit-downy-mildew/>

It's important to use fungicides with demonstrated performance in managing downy mildew and to treat preventatively for best control. The alphabetic list below was developed by Dr. Mary Hausbeck from fungicide performance in her Michigan trials in recent years. Fungicides should be alternated to mitigate the development of resistance.

- Elumin + chlorothalonil or mancozeb
- Omega (Orbus) + chlorothalonil or mancozeb
- Orondis Opti (chlorothalonil is part of the premix)
- Previcur Flex + chlorothalonil or mancozeb
- Ranman + chlorothalonil or mancozeb
- Zampro + chlorothalonil or mancozeb

For diagnostic and management support in Wisconsin, please consider contacting Dr. Amanda Gevens (gevens@wisc.edu) or our [Plant Disease Diagnostic Clinic \(pddc@wisc.edu\)](mailto:pddc@wisc.edu).


Vegetable Insect Update – Russell L. Groves, Professor and Associate Department Chairperson, UW-Madison, Department of Entomology, 608-262-3229 (office), (608) 698-2434 (cell), e-mail: rgroves@wisc.edu

Vegetable Entomology Webpage: <https://vegento.russell.wisc.edu/>

Potato virus Y and management. Important to note, aphid species that colonize potatoes (Green peach aphid and Potato aphid) were again detected in central Wisconsin fields over past week (July 5 – July 12) and also in seed potato fields in northern Wisconsin. The Aphid Suction Trap Network (valid through mid-June only) has indicated low numbers of potato-colonizing species have been detected flying in the landscape of the upper midwest. It is important to note, the data reported by the Suction Trap Network is 25 days late (June 15) and is only offered as evidence of low flight activity in mid June.

Initiate applications (of insecticides) after the appearance of colonizing aphids have become detected and established in the crop. An initial foliar application of a selected insecticide should be applied to the entire field and followed by a second foliar application of the same active ingredient one week later. For most products, only two successive applications are allowed per crop season for control of colonizing aphids before rotation to a new mode-of-action is suggested. Examples of exceptions to this include flonicamid ([Beleaf 50SG/Carbine 50WG®](#)) and imidacloprid ([AdmirePro®](#)), which can be applied over 3 successive foliar applications. Continue to scout field throughout the time interval of the application series and consider rotation to another effective aphicide mode of action if colonizing aphids persist in the crop.

All foliar applied aphicides should be applied with mild, penetrating surfactant and ensure the tank pH is not alkaline (pH < 7.0). Ground-application is also advised especially after row closure and when canopies begin to fall over in the late summer. Paraffinic oil applications can act as a mild penetrant, so not additional penetrating adjuvants are needed if paraffinic oils are applied with insecticides. As a reminder, all insecticide and paraffinic oil combinations should be delivered using application equipment that can ensure the most complete canopy coverage and using sufficient delivery volume.




Corn earworm – (<https://vegento.russell.wisc.edu/pests/corn-earworm/>).

Pheromone captures (as reported from the [Wisconsin Home Pest Survey](#)) now confirm adult moths of the corn earworm over the past week in select locations of the state. These captures are relatively low but numbers could increase in the coming week as conditions for moth flights were good in the past week ([Insect Forecast](#) site). These adults represent second generation, adult moths that are expected to continue flights into the region and silking sweet corn should be scouted for eggs and early instar larvae.

Once larvae enter the ear, they're protected from insecticides. Therefore, treatment timing is critical, and control must be targeted at the adults, eggs, and young larvae before they're protected. The amount of time for egg hatch varies with temperature, but during periods of hot summer days and nights, eggs can hatch within 24 hours after they're laid. Thus, each day's delay in treatment may reduce the number of clean ears by 10 to 15%.

Insecticides must be present on vulnerable silks when eggs hatch. The first application is the most critical and tank-mixes containing an ovicide (an egg killer) and a larvicide (eg. spinosad, spinetoram, chlorantraniliprole) are recommended. On sweet corn, treat every 5 to 7 days beginning when silks first appear and continuing until they are brown in color. Discontinue treatments 10 days before harvesting fresh market sweet corn as additional sprays will not improve product quality.




Moth Counts July 2-8, 2026

- Weekly total: 95
- Cumulative total: 95
- Traps reporting: 9
- Positive traps: 6
- Average moths per trap: 16
- Highest count: 36

County	Location	July 8
Columbia	Keyeser	
Columbia	Lodi	15
Columbia	Pardeeville	0
Dane	Sun Prairie N	
Dane	Sun Prairie S	
Dodge	Beaver Dam	8
Dodge	Leroy	0
Dodge	Watertown W	
Green Lake	Green Lake	0
Marathon	Wausau	1
Vernon	Coon Valley	
Walworth	Burlington	36
Walworth	Williams Bay	33
Wauzara	Hancock	2

https://datcp.wi.gov/Pages/Programs_Services/CornEarworm.aspx



<https://www.insectforecast.com/>

Langlade County Agriculture Research Station

Antigo Field Day 2026

Since 1975



Thursday
July 16
1:00 - 3:30

No Registration Fee

Presenters

UW Madison Agriculture Research Station Manager

- ★ **Kevin Gallenberg**
- **Welcome & Introduction**

UW Madison Department of Plant Pathology

- ★ **Amanda Gevens**
- **Potato Common Scab Research Update**

WI Seed Potato Certification Lab

- ★ **Brooke Babler**
- **Seed Potato Certification Updates**

UW Madison Department of Entomology

- ★ **Dr. Russ Groves**
- **PVY Full Season Management**

UW Madison Department of Agricultural & Life Sciences

- ★ **Maria Caraza-Harter**
- **UW-Madison Potato Breeding Program Updates**

UW Extension Regional Crops/Soils Educator

- ★ **Scott Reuss**
- **Cover Crops for Strong Rotations**



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON

Langlade County
Agriculture Research Station
N3689 Langlade Rd
Antigo, WI 54409



Wisconsin Potato Industry Board
Wisconsin Potato & Vegetable Growers Association
WPVGA Associate Division
Wisconsin Seed Potato Improvement Association
Mattek, J.W. & Sons, Inc.
Frito-Lay, Inc.
Baginski Farms
Schroeder Bros. Farms, Inc./Schroeder Farms, Ltd
Nutrien Ag Solution
Insight FS
Swiderski Equipment
Riesterer & Schnell
Langlade County
Rine Ridge Farms, Inc.
Bushman's Riverside Ranch, Inc.
Seidl Farms, Inc.
Sunnydale Farms, Inc.
Kohm & Spychalla, LLC



Following the Presentations - Food & Refreshments
Provided by Insight FS & WPVGA Associate Division at East City Park